

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 October 2002 (03.10.2002)

PCT

(10) International Publication Number
WO 02/076945 A1

(51) International Patent Classification⁷: C07D 207/48, 207/14, A61K 31/40

(21) International Application Number: PCT/US02/06475

(22) International Filing Date: 5 March 2002 (05.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/274,845 9 March 2001 (09.03.2001) US

(71) Applicant: ORTHO-MCNEIL PHARMACEUTICAL, INC. [US/US]; U.S. Route 202, Raritan, NJ 08869 (US).

(72) Inventors: GRECO, Michael N.,; 1634 Clearbrook Road, Lansdale, PA 19446 (US). MARYANOFF, Bruce E.,; 4029 Devonshire Drive, Forest Grove, PA 18922 (US). HAWKINS, Michael J.,; 318 Heckler Street, Ambler, PA 19002 (US). BOYD, Robert E.,; 3 Danbridge Drive, Horsham, PA 19044 (US).

(74) Agents: JOHNSON, Philip S., et al.; JOHNSON & JOHNSON, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).


(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/076945 A1

(54) Title: AMINOPYRROLIDINE SULFONAMIDES AS SERINE PROTEASE INHIBITORS

(57) Abstract: The present invention is directed to compounds useful as selective serine protease or dual-serine protease inhibitors, compositions thereof and methods for treating serine protease or dual-serine protease mediated disorders.

21052
#25

AMINOPYRROLIDINE SULFONAMIDES AS SERINE PROTEASE
INHIBITORS

CROSS REFERENCE TO RELATED APPLICATIONS

5

This application claims benefit of provisional application Serial Number 60/274,845, filed 9 March 2001, which is hereby incorporated by reference.

FIELD OF THE INVENTION

10

The present invention relates to certain novel compounds, methods for preparing the compounds, compositions, intermediates and derivatives thereof and for treating serine protease mediated disorders. More particularly, the aminopyrrolidine sulfonamide compounds of the present invention are selective
15 serine protease or dual-serine protease inhibitors of Factor Xa and tryptase useful for treating serine protease or dual-serine protease mediated disorders.

BACKGROUND OF THE INVENTION

20

Thrombotic disorders are a major cause of mortality in industrialized countries (Kaiser, Brigitte, Thrombin and Factor Xa Inhibitors, *Drugs of the Future*, 1998, 23(4), 423-436). Thrombin has been a target for the development of anticoagulation agents because it occupies a central position in the coagulation cascade (Kunitada, S., et al., Factor Xa Inhibitors, *Current*
25 *Pharmaceutical Design*, 1996, 2, 531-542). Since Factor Xa (FXa) is responsible for the formation of thrombin, a FXa inhibitor has become an alternative strategy to selectively prevent thrombin production and clot formation.

30

Like thrombin, FXa is a member of the serine protease superfamily. In the blood coagulation cascade, FXa links the intrinsic and extrinsic activation pathways for the production of thrombin. In the intrinsic pathway, Factor IXa converts Factor X to FXa in the presence of Factor VIIIa, Ca^{2+} and

phospholipid. In the extrinsic pathway, Factor VIIa converts Factor X to FXa in the presence of tissue factor. Once formed, FXa binds to Factor Va on phospholipid surfaces in the presence of Ca^{+2} ions to form the prothrombinase complex, which is responsible for converting prothrombin to thrombin.


- 5 Thrombin in turn converts fibrinogen to fibrin, which ultimately results in the production of a fibrin clot.

Potential advantages for FXa inhibitors as anticoagulants stem from the inhibition of thrombin formation rather than inhibition of its catalytic activity. For
10 example, it is expected that thrombin-induced platelet activation could still occur under FXa inhibition, thus minimizing bleeding risk. The thrombin/thrombomodulin complex downregulates thrombin production, thus functioning as an endogenous anticoagulant. It has been postulated that FXa inhibition would supply sufficient thrombin for this interaction, which might
15 minimize the "thrombotic rebound" effect observed in the clinical use of direct thrombin inhibitors.


A comprehensive review of FXa inhibitors has recently appeared (*Drugs of the Future* 1999, 24(7), 771-787).

20

PCT application WO 96/10022 to Faull, et. al., describes sulfonylpiperazine-derived FXa inhibitors of the formula:



PCT application WO 98/54164 to Tawada, et. al., describes
25 sulfonylpiperazine-derived FXa inhibitors of the formula:-




Accordingly, it is an object of the invention to provide aminopyrrolidine sulfonamide-derived compounds that are serine protease inhibitors; in particular, selective serine protease or dual-serine protease inhibitors of Factor Xa and tryptase. It is another object of the invention to provide a process for preparing aminopyrrolidine sulfonamide compounds, compositions, intermediates and derivatives thereof. It is a further object of the invention to provide methods for treating serine protease or dual-serine protease mediated disorders.


10

SUMMARY OF THE INVENTION

This invention is directed to aminopyrrolidine sulfonamide compounds selected from the group consisting of Formula (I) and Formula (II):



Formula (I)



Formula (II)

wherein

R_1 is selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{3-7} cycloalkyl, aryl, aryl(C_{1-8})alkyl, aryl(C_{2-8})alkenyl, heteroaryl(C_{1-8})alkyl, heteroaryl(C_{2-8})alkenyl and $R_4C(O)CH_2-$; wherein aryl and heteroaryl are optionally substituted with one to two substituents independently selected from R_4 ;

R_2 is selected from the group consisting of hydrogen, hydroxy, C_{1-8} alkoxy, aryloxy and aryl(C_{1-8})alkoxy; with the proviso that R_2 is bonded to the heterocyclyl ring by a single bond; alternatively, R_2 is oxo; with the proviso that R_2 is bonded to the heterocyclyl ring by a double bond;

R_3 is selected from the group consisting of aryl, aryl(C_{1-8})alkyl, aryl(C_{2-8})alkenyl, heteroaryl(C_{1-8})alkyl, heteroaryl(C_{2-8})alkenyl; wherein aryl and heteroaryl are optionally substituted with one to three substituents independently selected from the group consisting of halogen, C_{1-8} alkyl, C_{1-8} alkoxy, amino, (C_{1-4} alkyl)amino, di(C_{1-4} alkyl)amino, trihalo(C_{1-8})alkyl and trihalo(C_{1-8})alkoxy;

R_4 is selected from the group consisting of hydroxy, amino, C_{1-8} alkyl, (C_{1-4} alkyl)amino, di(C_{1-4} alkyl)amino, C_{1-8} alkoxy, carboxy, carboxy(C_{1-8})alkyl, carboxy(C_{1-8})alkoxy, (carboxy)amino, (carboxy(C_{1-4})alkyl)amino, (carboxyaryl)amino, (carboxyaryl(C_{1-4})alkyl)amino, (carboxy(C_{1-4})alkylaryl)amino, aryloxy, aryl(C_{1-8})alkoxy, (aryl)amino, (aryl(C_{1-4})alkyl)amino, (C_{1-4} alkylaryl)amino, (arylcarboxy)amino, di(aryl)amino, di(aryl(C_{1-4})alkyl)amino, C_{1-8} alkoxycarbonyl,

C₁₋₈alkoxycarbonyl(C₁₋₈)alkoxy, aminocarbonyl, (C₁₋₈alkyl)aminocarbonyl, (carboxy(C₁₋₈)alkyl)aminocarbonyl, (C₁₋₈alkoxycarbonyl(C₁₋₈)alkyl)aminocarbonyl and guanidino; and,

- 5 G is selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄alkyl, C₁₋₈alkoxy, aryl, aryloxy, aryl(C₁₋₈)alkyl, aryl(C₁₋₈)alkoxy, amino, carboxy, alkylaminocarbonyl, alkylcarbonylamino, trihalo(C₁₋₈)alkyl and trihalo(C₁₋₈)alkoxy;
- 10 and pharmaceutically acceptable salts thereof.

The aminopyrrolidine sulfonamide compounds of the present invention are selective serine protease or dual-serine protease inhibitors useful for treating serine protease mediated disorders. An embodiment of the invention

15 includes compounds that are selective or dual-inhibitors of Factor Xa and tryptase.

The present invention includes a method for preparing instant compounds, compositions, intermediates and derivatives thereof and a method

20 for treating selective serine protease or dual-serine protease mediated disorders.

DETAILED DESCRIPTION OF THE INVENTION

25 An embodiment of the instant compounds are those wherein, preferably, R₁ is selected from the group consisting of hydrogen, aryl(C₁₋₈)alkyl and heteroaryl(C₁₋₈)alkyl, wherein aryl, heteroaryl and the aryl and heteroaryl portion of arylalkyl and heteroarylalkyl are optionally substituted with a substituent selected from R₄. More preferably, R₁ is selected from the group

30 consisting of hydrogen, benzyl, phenethyl, phenylpropyl and benzofurylmethyl, wherein phenyl, the phenyl portion of benzyl and the benzofuryl portion of benzofurylmethyl are optionally substituted with a substituent selected from R₄. Most preferably, R₁ is selected from the group consisting of hydrogen, benzyl,

phenylpropyl and benzofurylmethyl, wherein phenyl, the phenyl portion of benzyl and the benzofuryl portion of benzofurylmethyl are optionally substituted with a substituent selected from R₄.

- 5 An embodiment of the instant compounds also include those wherein, preferably, R₂ is hydrogen.

- An embodiment of the instant compounds further includes those wherein, preferably, R₃ is aryl(C₂₋₈)alkenyl, wherein aryl is optionally substituted
10 with one to three substituents independently selected from halogen. More preferably, R₃ is independently selected from the group consisting of phenethenylene and phenylpropenylene, wherein phenyl is optionally substituted with one to three substituents independently selected from the group consisting of chlorine and fluorine. Most preferably, R₃ is
15 phenethenylene, wherein phenyl is substituted with one to three substituents selected from chlorine.

- Embodiments of the instant compounds include those wherein, preferably, R₄ is selected from the group consisting of hydroxy,
20 di(C₁₋₄alkyl)amino, C₁₋₈alkoxy, carboxy, carboxy(C₁₋₈)alkoxy, aryl(C₁₋₈)alkoxy, C₁₋₈alkoxycarbonyl, C₁₋₈alkoxycarbonyl(C₁₋₈)alkoxy, aminocarbonyl, (C₁₋₈alkyl)aminocarbonyl, (carboxy(C₁₋₈)alkyl)aminocarbonyl and C₁₋₈alkoxycarbonyl(C₁₋₈)alkyl)aminocarbonyl.

- 25 More preferably, R₄ is selected from the group consisting of hydroxy, carboxy, carboxy(C₁₋₈)alkoxy, C₁₋₈alkoxycarbonyl, C₁₋₈alkoxycarbonyl(C₁₋₈)alkoxy, aminocarbonyl, (carboxy(C₁₋₈)alkyl)aminocarbonyl and C₁₋₈alkoxycarbonyl(C₁₋₈)alkyl)aminocarbonyl.


30

 Most preferably, R₄ is selected from the group consisting of hydroxy, carboxy, carboxymethoxy, methoxycarbonyl, aminocarbonyl, (carboxymethylene)aminocarbonyl and

methoxycarbonylmethylene)aminocarbonyl.

- Embodiments of the instant compounds also include those wherein, preferably, G is selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄alkyl, C₁₋₈alkoxy, aryl, aryloxy, aryl(C₁₋₈)alkyl, aryl(C₁₋₈)alkoxy, amino and trihalo(C₁₋₈)alkyl. More preferably, G is hydrogen.

The compounds of the present invention are exemplified by a compound of the formula:




- 10 wherein R₄ is selected from:

Cpd	R ₄
2	4-OH;
3	4-CO ₂ CH ₃ ;
4	3-CO ₂ CH ₃ ;
5	4-CO ₂ H;
6	3-CO ₂ H;
7	3-OH;
8	3-OCH ₂ CO ₂ CH ₃ ;
9	4-CONH ₂ ;
10	4-CONHCH ₂ CO ₂ CH ₃ ;
11	3-OCH ₂ CO ₂ H;
12	4-CONHCH ₂ CO ₂ H;
13	3-CONHCH ₂ CO ₂ CH ₃ ;
14	3-CONHCH ₂ CO ₂ H;
15	3-CONH ₂ ;
24	4-NHC(=NH)NH ₂ ; or,
25	3-CO ₂ CH ₃ -4-OH;

and pharmaceutically acceptable salts thereof.

The compounds of the present invention are also exemplified by a compound of the formula:




wherein R₁ is:

Cpd	R ₁
1	PhCH ₂ ;
26	H; or,
27	2-benzofurylCH ₂

5 and pharmaceutically acceptable salts thereof.

The compounds of the present invention are further exemplified by a compound of the formula:



wherein R₁ and R₃ are dependently selected from:

Cpd	R ₁	R ₃
16	PhCH ₂	4-ClPh(CH ₂) ₂ ;
17	4-(PhCH ₂ O)PhCH ₂	4-ClPh(CH ₂) ₂ ;
18	4-[(CH ₃) ₂ N]PhCH ₂	4-ClPh(CH ₂) ₂ ;
19	4-(CH ₃ O)PhCH ₂	4-ClPh(CH ₂) ₂ ;
20	Ph(CH ₂) ₃	4-ClPh(CH ₂) ₂ ;
21	4-CO ₂ HPhCH ₂	4-ClPh(CH ₂) ₂ ;
22	4-[CH ₃ OC(O)]PhCH ₂	4-ClPh(CH ₂) ₂ ; or,
23	PhCH ₂	7-CH ₃ O-2-naphthalenyl;

10 and pharmaceutically acceptable salts thereof.

The compounds of the present invention may also be present in the form of a pharmaceutically acceptable salt. The pharmaceutically acceptable salt generally takes a form in which the basic nitrogen is protonated with an
5 inorganic or organic acid. Representative organic or inorganic acids include hydrochloric, hydrobromic, hydriodic, perchloric, sulfuric, nitric, phosphoric, acetic, propionic, glycolic, lactic, succinic, maleic, fumaric, malic, tartaric, citric, benzoic, mandelic, methanesulfonic, hydroxyethanesulfonic, benzenesulfonic, oxalic, pamoic, 2-naphthalenesulfonic, p-toluenesulfonic, cyclohexanesulfamic,
10 salicylic, saccharinic or trifluoroacetic.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds which are readily convertible *in vivo* into the
15 required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various disorders described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the subject. Conventional
20 procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

Where the compounds according to this invention have at least one
25 chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. Where the processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative
30 chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of

diastereomeric pairs by salt formation followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds
5 may be resolved using a chiral HPLC column. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention.

Resonance forms for compounds of the present invention include those
10 forms where an unsaturated bond may resonate between 2 or more atoms. For example, a guanidino group includes resonance forms represented by the formulae: -NH-C(=NH)-NH_2 or $\text{-N=C(NH}_2\text{)-NH}_2$. It is to be understood that all such resonance forms are encompassed within the scope of the present invention.

15 During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective
20 Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known in the art.

25 Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

30 The term "alkyl" refers to straight and branched-chain alkyl radical groups; similarly, alkenyl and alkynyl radicals include straight and branched chains having 2 to 8 carbon atoms or any number within this range; wherein

one or two double or triple bonds are formed in the chain between adjacent members. The term "alkoxy" refers to O-alkyl groups where alkyl is as defined supra. The term "cycloalkyl" refers to a cyclic alkyl ring of five to seven carbon atom members. Examples of such cyclic alkyl rings include pentyl, hexyl or
5 heptyl.

The term "heterocyclyl" refers to a saturated or partially unsaturated ring having five members of which at least one member is a N, O or S atom and which optionally contains one additional O atom or one, two or three additional
10 N atoms, a saturated or partially unsaturated ring having six members of which one, two or three members are a N atom, a saturated or partially unsaturated bicyclic ring having nine members of which at least one member is a N, O or S atom and which optionally contains one, two or three additional N atoms or a
15 saturated or partially unsaturated bicyclic ring having ten members of which one, two or three members are a N atom. Examples include, and are not limited to, pyrrolinyl, pyrrolidinyl, 1,3-dioxolanyl, imidazoliny, imidazolidinyl, pyrazoliny, pyrazolidinyl, piperidinyl, morpholinyl or piperazinyl.

The term "aryl" refers to a single aromatic ring of six carbon members or
20 a bicyclic aromatic ring of ten carbon members. Examples of such aryl rings include phenyl and naphthyl.

The term "heteroaryl" refers to an aromatic monocyclic ring system containing five members of which at least one member is a N, O or S atom and
25 which optionally contains one, two or three additional N atoms; an aromatic monocyclic ring having six members of which one, two or three members are a N atom; an aromatic bicyclic ring having nine members of which at least one member is a N, O or S atom and which optionally contains one, two or three additional N atoms; or, an aromatic bicyclic ring having ten members of which
30 one, two or three members are a N atom. Examples include, and are not limited to, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, indazolyl, benzo[b]furyl, benzo[b]thienyl, quinolinyl, isoquinolinyl or quinazolinyl.

The term "carbonyl" as used herein refers to the organic radical linking group: R-C(O)-R having a single carbon atom; the term "carboxy" as used herein refers to the organic radical terminal group: R-C(O)OH.

5

The term "halogen" or "halo" shall include iodine, bromine, chlorine and fluorine.


Whenever the term "alkyl" or "aryl" or either of their prefix roots appear in a name of a substituent (e.g., aryl(C₁-C₄)alkyl, di(C₁-C₄ alkyl)amino) it shall be interpreted as including those limitations given above for "alkyl" and "aryl." Designated numbers of carbon atoms (e.g., C₁-C₆) shall refer independently to the number of carbon atoms in an alkyl or cycloalkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

15

The term "aryl(C₁₋₈)alkyl" or the coextensive term "arylalkyl" means an alkyl group substituted at the terminal carbon with an aryl group (e.g., benzyl, phenethyl). Similarly, the term "heteroaryl(C₁₋₈)alkyl" or the coextensive term "heteroarylalkyl" means an alkyl group substituted at the terminal carbon with a heteroaryl group. The term "aryl(C₁₋₈)alkoxy" or the coextensive term "arylalkoxy" indicates an alkoxy group substituted at the terminal carbon with an aryl group (e.g., benzyloxy).

When a particular group is "substituted" (e.g., Phe, aryl, heteroaryl), that group may have one or more substituents, preferably from one to five substituents, more preferably from one to three substituents, most preferably from one to two substituents, independently selected from the list of substituents.

Under standard nomenclature rules used throughout this disclosure, the terminal portion of the designated side chain is described first followed by the adjacent functionality toward the point of attachment. Thus, for example, a "phenylC₁₋₆alkylamidoC₁₋₆alkyl" substituent refers to a group of the formula:



It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein.

The aminopyrrolidine sulfonamide-derived compounds of the present invention are selective serine protease or dual-serine protease inhibitors; in particular, selective or dual-inhibitors of Factor Xa and tryptase useful for treating selective serine protease or dual-serine protease mediated disorders.

Embodiments of the method of the present invention include a method for treating or ameliorating a serine protease or dual-serine protease mediated disorder in a subject in need thereof comprising administering to the subject a therapeutically effective amount of an instant compound or pharmaceutical composition thereof. The therapeutically effective amount of the compounds selected from the group consisting of Formula (I) and Formula (II) exemplified in such a method is from about 0.001 mg/kg/day to about 300 mg/kg/day.

Embodiments of the present invention include the use of a compound selected from the group consisting of Formula (I) and Formula (II) for the preparation of a medicament for treating or ameliorating a serine protease or dual-serine protease mediated disorder in a subject in need thereof.

In accordance with the methods of the present invention, an individual compound of the present invention or a pharmaceutical composition thereof

can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted
5 accordingly.

Embodiments of the present method include a compound or pharmaceutical composition thereof advantageously co-administered in combination with other agents for treating or ameliorating a serine protease or
10 dual-serine protease mediated disorder. For example, in the treatment of thrombosis, a compound selected from the group consisting of Formula (I) and Formula (II) or pharmaceutical composition thereof may be used in combination with other agents.

15 The combination product comprises co-administration of a compound selected from the group consisting of Formula (I) and Formula (II) or pharmaceutical composition thereof and an additional agent for treating or ameliorating a serine protease or dual-serine protease mediated disorder, the sequential administration of a compound selected from the group consisting of
20 Formula (I) and Formula (II) or pharmaceutical composition thereof and an additional agent for treating or ameliorating a serine protease or dual-serine protease mediated disorder, administration of a pharmaceutical composition containing a compound selected from the group consisting of Formula (I) and Formula (II) or pharmaceutical composition thereof and an additional agent for
25 treating or ameliorating a serine protease or dual-serine protease mediated disorder or the essentially simultaneous administration of a separate pharmaceutical composition containing a compound selected from the group consisting of Formula (I) and Formula (II) or pharmaceutical composition thereof and a separate pharmaceutical composition containing an additional
30 agent for treating or ameliorating a serine protease or dual-serine protease mediated disorder.

The term "subject" as used herein, refers to an animal, preferably a

mammal, most preferably a human, who has been the object of treatment, observation or experiment.

5 The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human, that is being sought by a researcher, veterinarian, medical doctor, or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

10 The ubiquitous nature of the Factor Xa and tryptase isoforms and their important roles in physiology provide incentive to produce highly selective Factor Xa and tryptase inhibitors. Given the evidence demonstrating linkage of certain isoforms to disease states, it is reasonable to assume that inhibitory compounds that are selective to a serine protease isoform or to a Factor Xa
15 isoform relative to the a tryptase isoform and other serine proteases or dual-serine protease are superior therapeutic agents. Such compounds should demonstrate greater efficacy and lower toxicity by virtue of their specificity. Accordingly, it will be appreciated by one skilled in the art that a compound selected from the group consisting of Formula (I) and Formula (II) is
20 therapeutically effective for certain serine protease or dual-serine protease mediated disorders based on the modulation of the disorder by selective serine protease or dual-serine protease inhibition. The usefulness of a compound selected from the group consisting of Formula (I) and Formula (II) as a
25 selective serine protease or dual-serine protease inhibitor can be determined according to the methods disclosed herein and the scope of such use includes use in one or more serine protease or dual-serine protease mediated disorders.

30 More particularly, the term "serine protease or dual-serine protease mediated disorders" includes, and is not limited to, thrombotic disorders, arterial thrombosis, venous thrombosis, restenosis, hypertension, heart failure, arrhythmia, myocardial infarction, acute myocardial infarction, reocclusion following thrombolytic therapy, reocclusion following angioplasty, inflammation, angina, unstable angina, stroke, atherosclerosis, ischemic conditions,

neurodegenerative disorders (associated with thrombotic or ischemic conditions), asthma and inflammatory bowel syndrome. Certain of the instant compounds are also useful as antithrombotics and anticoagulation agents in conjunction with fibrinolytic therapy (e.g., t-PA or streptokinase). The utility of
5 the compounds to treat serine protease or dual-serine protease mediated disorders can be determined according to the procedures described herein.

Additionally, compounds of the present invention (such as Compound 27) may be useful in treating a chronic neurodegenerative disorder (such as
10 Alzheimer's disease) (tested for activity using methodology similar to that described in Ermolieff, J., et al, *Proteolytic Activation of Recombinant Pro-memapsin 2 (Pro- β -secretase) Studied with New Fluorogenic Substrates, Biochemistry, 2000, 39, 12450-12456).*

15 The present invention further provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier.

As used herein, the term "composition" is intended to encompass a
20 product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts. Accordingly, pharmaceutical compositions containing the compounds of the present invention as the active ingredient as well as methods of preparing the instant compounds are also part
25 of the present invention.

To prepare the pharmaceutical compositions of this invention, one or more compounds of Formula (I), Formula (II) or salt thereof as the active ingredient, is intimately admixed with a pharmaceutical carrier according to
30 conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration (e.g. oral or parenteral). Suitable pharmaceutically acceptable carriers are well known in the art. Descriptions of some of these

pharmaceutically acceptable carriers may be found in The Handbook of Pharmaceutical Excipients, published by the American Pharmaceutical Association and the Pharmaceutical Society of Great Britain.

- 5 Methods of formulating pharmaceutical compositions have been described in numerous publications such as Pharmaceutical Dosage Forms: Tablets, Second Edition, Revised and Expanded, Volumes 1-3, edited by Lieberman, et al.; Pharmaceutical Dosage Forms: Parenteral Medications, Volumes 1-2, edited by Avis, et al.; and Pharmaceutical Dosage Forms: Disperse Systems, Volumes 1-2, edited by Lieberman, et al.; published by Marcel Dekker, Inc.
- 10

 In preparing a pharmaceutical composition of the present invention in liquid dosage form for oral, topical and parenteral administration, any of the
15 usual pharmaceutical media or excipients may be employed. Thus, for liquid dosage forms, such as suspensions (i.e. colloids, emulsions and dispersions) and solutions, suitable carriers and additives include but are not limited to pharmaceutically acceptable wetting agents, dispersants, flocculation agents, thickeners, pH control agents (i.e. buffers), osmotic agents, coloring agents,
20 flavors, fragrances, preservatives (i.e. to control microbial growth, etc.) and a liquid vehicle may be employed. Not all of the components listed above will be required for each liquid dosage form:

 In solid oral preparations such as, for example, powders, granules,
25 capsules, caplets, gelcaps, pills and tablets (each including immediate release, timed release and sustained release formulations), suitable carriers and additives include but are not limited to diluents, granulating agents, lubricants, binders, glidants, disintegrating agents and the like. Because of their ease of administration, tablets and capsules represent the most advantageous oral
30 dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated, gelatin coated, film coated or enteric coated by standard techniques.

The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described above. The pharmaceutical compositions herein will contain, per unit dosage unit, e.g.,
5 tablet, capsule, powder, injection, suppository, teaspoonful and the like, of from about 0.01 mg to about 300 mg (preferably, from about 0.1 mg to about 100 mg; and, more preferably, from about 1 mg to about 30 mg) and may be given at a dosage of from about 0.01 mg/kg/day to about 300 mg/kg/day (preferably, from about 0.1 mg/kg/day to about 100 mg/kg/day; and, more preferably, from
10 about 1 mg/kg/day to about 30 mg/kg/day). Preferably, in the method for treating thrombotic disorders described in the present invention and using any of the compounds as defined herein, the dosage form will contain a pharmaceutically acceptable carrier containing between about 0.01 mg and 100 mg; and, more preferably, between about 5 mg and 50 mg of the compound; and,
15 may be constituted into any form suitable for the mode of administration selected. The dosages, however, may be varied depending upon the requirement of the subjects, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

20

Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, lozenges, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories for administration by oral, intranasal, sublingual,
25 *intraocular, transdermal, parenteral, rectal, vaginal, inhalation or insufflation means. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection.*

30

For preparing solid pharmaceutical compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as diluents, binders, adhesives,

disintegrants, lubricants, antiadherents and glidants. Suitable diluents include, but are not limited to, starch (i.e. corn, wheat, or potato starch, which may be hydrolyzed), lactose (granulated, spray dried or anhydrous), sucrose, sucrose-based diluents (confectioner's sugar; sucrose plus about 7 to 10 weight percent invert sugar; sucrose plus about 3 weight percent modified dextrins; sucrose plus invert sugar, about 4 weight percent invert sugar, about 0.1 to 0.2 weight percent cornstarch and magnesium stearate), dextrose, inositol, mannitol, sorbitol, microcrystalline cellulose (i.e. AVICEL™ microcrystalline cellulose available from FMC Corp.), dicalcium phosphate, calcium sulfate dihydrate, calcium lactate trihydrate and the like. Suitable binders and adhesives include, but are not limited to acacia gum, guar gum, tragacanth gum, sucrose, gelatin, glucose, starch, and cellulose (i.e. methylcellulose, sodium carboxymethylcellulose, ethylcellulose, hydroxypropylmethylcellulose, hydroxypropylcellulose, and the like), water soluble or dispersible binders (i.e. alginic acid and salts thereof, magnesium aluminum silicate, hydroxyethylcellulose [i.e. TYLOSE™ available from Hoechst Celanese], polyethylene glycol, polysaccharide acids, bentonites, polyvinylpyrrolidone, polymethacrylates and pregelatinized starch) and the like. Suitable disintegrants include, but are not limited to, starches (corn, potato, etc.), sodium starch glycolates, pregelatinized starches, clays (magnesium aluminum silicate), celluloses (such as crosslinked sodium carboxymethylcellulose and microcrystalline cellulose), alginates, pregelatinized starches (i.e. corn starch, etc.), gums (i.e. agar, guar, locust bean, karaya, pectin, and tragacanth gum), cross-linked polyvinylpyrrolidone and the like. Suitable lubricants and antiadherents include, but are not limited to, stearates (magnesium, calcium and sodium), stearic acid, talc waxes, stearowet, boric acid, sodium chloride, DL-leucine, carbowax 4000, carbowax 6000, sodium oleate, sodium benzoate, sodium acetate, sodium lauryl sulfate, magnesium lauryl sulfate and the like. Suitable glidants include, but are not limited to, talc, cornstarch, silica (i.e. CAB-O-SIL™ silica available from Cabot, SYLOID™ silica available from W.R. Grace/Davison, and AEROSIL™ silica available from Degussa) and the like. Sweeteners and flavorants may be added to chewable solid dosage forms to improve the palatability of the oral dosage form. Additionally, colorants and

coatings may be added or applied to the solid dosage form for ease of identification of the drug or for aesthetic purposes. These carriers are formulated with the pharmaceutical active to provide an accurate, appropriate dose of the pharmaceutical active with a therapeutic release profile.

5

Generally these carriers are mixed with the pharmaceutical active to form a solid preformulation composition containing a homogeneous mixture of the pharmaceutical active of the present invention, or a pharmaceutically acceptable salt thereof. Generally the preformulation will be formed by one of
10 three common methods: (a) wet granulation, (b) dry granulation and (c) dry blending. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This
15 solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from about 0.1 mg to about 500 mg of the active ingredient of the present invention. The tablets or pills containing the novel compositions may also be formulated in multilayer tablets or pills to provide a sustained or provide dual-release products. For example, a dual
20 release tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer, which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be
25 used for such enteric layers or coatings, such materials including a number of polymeric materials such as shellac, cellulose acetate such as cellulose acetate phthalate, polyvinyl acetate phthalate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, methacrylate and ethylacrylate copolymers, methacrylate and methyl methacrylate copolymers
30 and the like. Sustained release tablets may also be made by film coating or wet granulation using slightly soluble or insoluble substances in solution (which for a wet granulation acts as the binding agents) or low melting solids a molten form (which in a wet granulation may incorporate the active ingredient). These

materials include natural and synthetic polymers waxes, hydrogenated oils, fatty acids and alcohols (i.e. beeswax, carnauba wax, cetyl alcohol, cetylstearyl alcohol, and the like), esters of fatty acids metallic soaps, and other acceptable materials that can be used to granulate, coat, entrap or otherwise limit the solubility of an active ingredient to achieve a prolonged or sustained release product.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, but are not limited to aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable suspending agents for aqueous suspensions, include synthetic and natural gums such as, acacia, agar, alginate (i.e. propylene alginate, sodium alginate and the like), guar, karaya, locust bean, pectin, tragacanth, and xanthan gum, cellulosics such as sodium carboxymethylcellulose, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose and hydroxypropyl methylcellulose, and combinations thereof, synthetic polymers such as polyvinyl pyrrolidone, carbomer (i.e. carboxypolymethylene), and polyethylene glycol; clays such as bentonite, hectorite, attapulgite or sepiolite; and other pharmaceutically acceptable suspending agents such as lecithin, gelatin or the like. Suitable surfactants include but are not limited to sodium docusate, sodium lauryl sulfate, polysorbate, octoxynol-9, nonoxynol-10, polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 80, polyoxamer 188, polyoxamer 235 and combinations thereof. Suitable deflocculating or dispersing agent include pharmaceutical grade lecithins. Suitable flocculating agent include but are not limited to simple neutral electrolytes (i.e. sodium chloride, potassium, chloride, and the like), highly charged insoluble polymers and polyelectrolyte species, water soluble divalent or trivalent ions (i.e. calcium salts, alums or sulfates, citrates and phosphates (which can be used jointly in formulations as pH buffers and flocculating agents). Suitable preservatives include but are not limited to parabens (i.e. methyl, ethyl, *n*-propyl and *n*-butyl), sorbic acid,

thimerosal, quaternary ammonium salts, benzyl alcohol, benzoic acid, chlorhexidine gluconate, phenylethanol and the like. There are many liquid vehicles that may be used in liquid pharmaceutical dosage forms, however, the liquid vehicle that is used in a particular dosage form must be compatible with the suspending agent(s). For example, nonpolar liquid vehicles such as fatty esters and oils liquid vehicles are best used with suspending agents such as low HLB (Hydrophile-Lipophile Balance) surfactants, stearylalkonium hectorite, water insoluble resins, water insoluble film forming polymers and the like. Conversely, polar liquids such as water, alcohols, polyols and glycols are best used with suspending agents such as higher HLB surfactants, clays silicates, gums, water soluble cellulose, water soluble polymers and the like. For parenteral administration, sterile suspensions and solutions are desired. Liquid forms useful for parenteral administration include sterile solutions, emulsions and suspensions. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

Furthermore, compounds of the present invention can be administered in an intranasal dosage form via topical use of suitable intranasal vehicles or via transdermal skin patches, the composition of which are well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the administration of a therapeutic dose will, of course, be continuous rather than intermittent throughout the dosage regimen.

Compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, multilamellar vesicles and the like. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, phosphatidylcholines and the like.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include, but are

- not limited to polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxy-ethylaspartamidephenol, or polyethyl eneoxydepolylysine substituted with palmitoyl residue. Furthermore, the compounds of the present invention may be coupled to a class of
- 5 biodegradable polymers useful in achieving controlled release of a drug, for example, to homopolymers and copolymers (which means polymers containing two or more chemically distinguishable repeating units) of lactide (which includes lactic acid d-, l- and meso lactide), glycolide (including glycolic acid), ϵ -caprolactone, p-dioxanone (1,4-dioxan-2-one), trimethylene carbonate (1,3-
- 10 dioxan-2-one), alkyl derivatives of trimethylene carbonate, δ -valerolactone, β -butyrolactone, γ -butyrolactone, ϵ -decalactone, hydroxybutyrate, hydroxyvalerate, 1,4-dioxepan-2-one (including its dimer 1,5,8,12-tetraoxacyclotetradecane-7,14-dione), 1,5-dioxepan-2-one, 6,6-dimethyl-1,4-dioxan-2-one, polyorthoesters, polyacetals, polydihydropyrans,
- 15 polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels and blends thereof.

Compounds of this invention may be administered in any of the foregoing compositions and dosage regimens or by means of those compositions and

20 dosage regimens established in the art whenever treatment of serine protease or dual-serine protease mediated disorders is required for a subject in need thereof.

The daily dose of a pharmaceutical composition of the present invention may be varied over a wide range from about 0.7 mg to about 21,000 mg per 70

25 kilogram (kg) adult human per day; preferably in the range of from about 0.7 mg to about 7,000 mg per adult human per day; and, more preferably, in the range of from about 0.7 mg to about 2,100 mg per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200,

30 250 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the subject to be treated. A therapeutically effective amount of the drug is ordinarily supplied at a dosage level of from about 0.01 mg/kg to about 300 mg/kg of body weight per day. Preferably, the range is from about 0.1 mg/kg

to about 100 mg/kg of body weight per day; and, most preferably, from about 1 mg/kg to about 30 mg/kg of body weight per day. Advantageously, compounds of the present invention may be administered in a single daily dose or the total daily dosage may be administered in divided doses of two, three or four times daily.

Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, and the advancement of the disease condition. In addition, factors associated with the particular subject being treated, including subject age, weight, diet and time of administration, will result in the need to adjust the dose to an appropriate therapeutic level.

In the examples and throughout this application, the following abbreviations have the meanings recited hereinafter:

Boc	t-Butoxycarbonyl
Cpd or Cpmd	Compound
DCC	1,3-Dicyclohexylcarbodiimide
DIPEA	Diisopropylethylamine
DMF	N, N-Dimethylformamide
EtOAc	Ethyl acetate
h	Hour
KHSO ₄	Potassium bisulfate
MeOH	Methanol
min	Minute
mL	Milliliter
NaBH ₄	Sodium borohydride
Na ₂ SO ₄	Sodium sulfate
NaHCO ₃	Sodium bicarbonate
rt	room temperature
TFA	Trifluoroacetic acid

General Synthetic Examples

Representative compounds of the present invention can be synthesized in accordance with the general synthetic methods described below and are

illustrated more particularly in the schemes that follow. Since the schemes are illustrations, the invention should not be construed as being limited by the chemical reactions and conditions expressed. The preparation of the various starting materials used in the schemes is well within the skill of persons versed
5 in the art.

The following scheme describes general synthetic methods whereby intermediate and target compounds of the present invention may be prepared. Additional representative compounds of the present invention can be
10 synthesized using the intermediates prepared in accordance with the schemes and other materials, compounds and reagents known to those skilled in the art.

General Synthetic Examples

Representative compounds of the present invention can be synthesized
15 in accordance with the general synthetic methods described below and are illustrated more particularly in the schemes that follow. Since the schemes are illustrations, the invention should not be construed as being limited by the chemical reactions and conditions expressed. The preparation of the various starting materials used in the schemes is well within the skill of persons versed
20 in the art.

The following scheme describes general synthetic methods whereby intermediate and target compounds of the present invention may be prepared. Additional representative compounds of the present invention can be
25 synthesized using the intermediates prepared in accordance with the schemes and other materials, compounds and reagents known to those skilled in the art.

Scheme A illustrates a general synthetic route for compounds of Formula (I).
30

In a modification of a known procedure (*Syn. Comm.* **1992**, 22(19) 2357), the 3-aminopyrrolidine Compound A1 was treated with an aldehyde in a solvent such as toluene and the mixture was heated at reflux with water removal by a


Dean-Stark apparatus. The required pyrrolidine intermediate for preparing compounds wherein R_2 is oxo may be made according to PCT application WO 98/05336. The required intermediate for compounds wherein R_1 is H may be made by using a commercially available 3-butyloxycarbonylamino pyrrolidine or other known analog. Upon cooling to rt, the solution was treated with a protecting group reagent such as di-*tert*-butyl dicarbonate. The intermediate imine was dissolved in an anhydrous solvent such as methanol and treated with a reducing agent such as NaBH_4 to afford an amine Compound **A2**. Treatment of Compound **A2** with an acid chloride such as 4-nitrobenzoyl chloride in a solvent such as CH_2Cl_2 followed by solvent removal and treatment with TFA afforded Compound **A3**.

Sulfonylation of Compound **A3** can be carried out by treating Compound **A3** with a sulfonyl halide in a solvent such as CH_2Cl_2 in the presence of an amine base such as DIPEA to afford Compound **A4**.

Treatment of Compound **A4** with a reducing agent such as SnCl_2 in the presence of a mineral acid such as HCl in an alcoholic solvent such as MeOH afforded Compound **A5**.

Guanylation of Compound **A5** was carried out by treating Compound **A5** with a guanylating agent such *N,N*-bis-*tert*-butoxycarbonylthiourea in the presence of HgCl_2 , an amine base such as triethylamine in a solvent such as DMF to afford Compound **A6**. Treatment of Compound **A6** with trifluoroacetic acid yielded the target compound of **Formula (I)**.

Scheme A



Scheme B illustrates the general synthetic route for compounds of Formula (II).

5


The intermediate Compound A2 (Scheme A) was treated with a sulfonyl halide in a solvent such as CH_2Cl_2 in the presence of an amine base such as DIPEA to afford Compound B1. Treatment of Compound B1 with TFA in a solvent such as CH_2Cl_2 gave Compound B2. A mixture of Compound B2, an aromatic carboxylic acid chloride such as 4-nitrobenzoyl chloride and a base such as DIPEA in a solvent such as CH_2Cl_2 were stirred at rt to afford

Compound **B3**.

Treatment of Compound **B3** with a reducing agent such as SnCl_2 in the presence of a mineral acid such as HCl in an alcoholic solvent such as MeOH
5 afforded Compound **B4**.

Guanylation of Compound **B4** was carried out by treating Compound **B4** with a guanylation agent such as *N,N'*-bis-*tert*-butoxycarbonythiourea in the presence of HgCl_2 and an amine base such as triethylamine in a solvent such
10 as DMF to afford Compound **B5**. Treatment of Compound **B5** with trifluoroacetic acid in a solvent such as CH_2Cl_2 yielded the target compound of Formula (II).

Scheme B

Specific Synthetic Examples

- 5 Specific compounds which are representative of this invention were prepared as per the following examples and reaction sequences; the examples and the diagrams depicting the reaction sequences are offered by way of illustration, to aid in the understanding of the invention and should not be construed to limit in

any way the invention set forth in the claims which follow thereafter. The depicted intermediates may also be used in subsequent examples to produce additional compounds of the present invention. No attempt has been made to optimize the yields obtained in any of the reactions. One skilled in the art would
5 know how to increase such yields through routine variations in reaction times, temperatures, solvents and/or reagents.

All chemicals were obtained from commercial suppliers and used without further purification. ^1H and ^{13}C NMR spectra were recorded on a Bruker AC
10 300B (300 MHz proton) or a Bruker AM-400 (400 MHz proton) spectrometer with Me_4Si as an internal standard (s = singlet, d = doublet, t = triplet, br = broad). APCI-MS and ES-MS were recorded on a VG Platform II mass spectrometer. TLC was performed with Whatman 250- μm silica gel plates. Preparative TLC was performed with Analtech 1000- μm silica gel GF plates.
15 Flash column chromatography was conducted with flash column silica gel (40-63 μm) and column chromatography was conducted with standard silica gel. HPLC separations were carried out on three Waters PrepPak[®] Cartridges (25 x 100 mm, Bondapak[®] C18, 15-20 μm , 125 Å) connected in series; detection was at 254 nm on a Waters 486 UV detector. Analytical HPLC was carried out on a
20 Supelcosil ABZ+PLUS column (5 cm x 2.1 mm), with detection at 254 nm on a Hewlett Packard 1100 UV detector. Microanalysis was performed by Robertson Microlit Laboratories, Inc.

Representative Chemical Abstracts Service (CAS) Index-like names for
25 the compounds of the present invention were derived using the ACD/LABS SOFTWARE[™] Index Name Pro Version 4.5 nomenclature software program provided by Advanced Chemistry Development, Inc., Toronto, Ontario, Canada.

Example 1

30 4-[[[1-[[*(E)*-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]-benzoic acid (Compound 5)

To a solution of Compound 1A (1.4g, 16.0 mMol) in toluene (65 mL) was added

methyl 4-formyl benzoate (2.6g, 16.0 mMol). The reaction was refluxed (Dean-Stark trap) until water distillation ended (~30 min). The solution was cooled to rt and treated with di-*tert*-butyl dicarbonate (3.1 g, 16.0 mMol) in one portion. After stirring overnight, the reaction was concentrated to yield Compound **1B** as a yellow-brown oil: ^1H NMR (CDCl_3) δ 1.37-1.58 (m, 9H), 1.71-2.22 (overlapping m, 3H), 3.32-3.73 (overlapping m, 4H), 3.88-4.09 (overlapping m, 4H), 7.71 (d, 2H), 8.09 (d, 2H), 8.37 (s, 1H) ppm.

To a solution of Compound **1B** at 0°C in 50 mL of MeOH was added excess NaBH_4 (0.8 g, 21.1 mMol). The ice bath was removed and the solution was warmed to rt. To the solution was added 3 mL of acetone, and the solution was concentrated to yield a gummy solid which was partitioned between EtOAc and brine. The layers were separated and the aqueous layer was extracted with EtOAc. The combined extracts were filtered and dried over Na_2SO_4 . The solution was filtered through Celite and concentrated to yield Compound **1C** as an orange oil: 5.5g; HPLC: 2.42 min, 100%; ES-MS 335(MH^+); ^1H NMR (CDCl_3) δ 1.48 (s, 9H), 1.66-1.81 (m, 1H), 1.98-2.11 (m, 1H), 3.04-3.22 (m, 1H), 3.26-3.62 (overlapping m, 4H), 3.87 (s, 2H), 3.92 (s, 3H), 7.4 (d, 2H), 8.0 (d, 2H).

To a solution of Compound **1C** (1.1 g, 3.4 mMol) in CH_2Cl_2 (30 mL) was added DIPEA (0.6 mL, 3.4 mMol) followed by *p*-nitrobenzoyl chloride (0.6 g, 3.4 mMol). After 40 min, the reaction was concentrated and the residue was treated with 25 mL of TFA and stirred for 25 min. The solution was concentrated to yield Compound **1D** as an orange oil: HPLC 2.55 min, 91%; ES-MS 384 (MH^+).


To a solution of Compound **1D** in CH_2Cl_2 (30 mL) was added DIPEA (6.5 mL, 37.3 mMol) followed by 0.9 g, 3.7 mMol of *p*-chlorolstyrylsulfonyl chloride (prepared as in WO 96/10022) and stirred for 30 min. The reaction solution was washed sequentially with 2X 1N KHSO_4 , 2X sat. NaHCO_3 , 1X brine, and filtered. The filtrate was dried over Na_2SO_4 , filtered through Celite, and concentrated to yield Compound **1E** as a brown foam: 2.0g; HPLC 4.04 min, 85%.

To a suspension of Compound 1E (1.8 g, 3.1 mMol) in MeOH (40 mL) was added a solution of SnCl₂ (2.9 g, 15.4 mMol) in concentrated HCl (10 mL) and refluxed for 1.5h. The solution was concentrated, made basic (blue litmus) 1N NaOH and extracted twice with EtOAc. The combined extracts were washed with brine, dried over Na₂SO₄, filtered and concentrated to yield 1.5 g of Compound 1F as a yellow glass;; HPLC 3.38 min, 81%.

To a solution of Compound 1F (1.6 g, 2.9 mMol) in DMF (30 mL) was added (1.2 g, 4.4 mMol) of *N,N'*-bis-*tert*-butoxycarbonylthiourea (*Syn. Comm.* 1993, 23(10), 1443.) followed by TEA (1.4 mL, 9.7 mMol). To the solution was added 1.2g (4.4 mMol) of HgCl₂. After 5 h, the black reaction mixture was diluted with EtOAc (150 mL) and filtered through Celite. The resulting light orange solution was washed sequentially with water and brine, filtered, dried over Na₂SO₄ and concentrated. The residue was purified by flash column chromatography (silica gel, CH₂Cl₂:MeOH; 100% -> 99:1) to yield Compound 1G as a white solid: (0.9g, 38.5%); HPLC 99% 4.48 min; ES-MS 796 (MH⁺).

To a solution of Compound 1G (0.1 g, 0.1 mMol) in 1,4-dioxane (18 mL) was added a solution of LiOH·H₂O (16 mg, 0.4 mMol) in H₂O (2 mL) and stirred overnight. The reaction solution was concentrated, acidified with excess 1N KHSO₄, and extracted three times with EtOAc. The combined extracts were washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash column chromatography (silica gel, 98:2 CH₂Cl₂:MeOH) to afford Compound 1H as a white powder: (48 mg, 51%); HPLC 4.25 min; 99%.

A solution of Compound 1H (48 mg, 0.06 mMol) was dissolved in 2 mL of TFA and stirred for 30 min. The solution was concentrated to yield a clear oil which was triturated 2X Et₂O and dried under vacuum at rt to yield Compound 5 as a white solid: (37.0 mg, 85%); HPLC 2.76 min, 100%. ES-MS 582 (MH⁺); ¹H NMR (DMSO-d₆) δ 1.88-2.12 (broad s, 2H), 2.92-3.53 (overlapping m, 4H), 4.38-4.58 (overlapping m, 3H), 7.07-7.98 (m, 18H), 9.98 (broad s, 1H).



Using the procedure of Example 1 and the appropriate reagents and starting materials known to those skilled in the art, other compounds of the present invention may be prepared including, but not limited to:

Cpd	Name	ES-MS m/z (MH ⁺)
1	<i>N</i> -[1-[[<i>(E)</i> -2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl]-4-[(diaminomethylene)amino]- <i>N</i> -(phenylmethyl)-benzamide	538
2	<i>N</i> -[1-[[<i>(E)</i> -2-(4-chlorophenyl)ethenyl]sulfonyl]-3-	554

	pyrrolidinyl]-4-[(diaminomethylene)amino]-N-[(4-hydroxyphenyl)methyl]-benzamide	
3	4-[[[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]-benzoic acid methyl ester	596
4	3-[[[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]-benzoic acid methyl ester	596
6	3-[[[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]-benzoic acid	582
7	<i>N</i> -[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl]-4-[(diaminomethylene)amino]- <i>N</i> -[(3-hydroxyphenyl)methyl]-benzamide	554
8	[3-[[[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]phenoxy]-acetic acid methyl ester	626
9	<i>N</i> -[[4-(aminocarbonyl)phenyl]methyl]- <i>N</i> -[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl]-4-[(diaminomethylene)amino]-benzamide,	581
10	[4-[[[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]benzoyl]amino]-acetic acid methyl ester	653
11	[3-[[[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]phenoxy]-acetic acid	612
12	[4-[[[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]benzoyl]amino]-acetic acid	639
13	[3-[[[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]benzoyl]amino]-acetic acid methyl ester	653
14	[3-[[[1-[(<i>E</i>)-2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]benzoyl]amino]-acetic acid	639

15	3-[[[1-[[<i>(E)</i> -2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]-benzamide	581
24	<i>N</i> -[[4-[(aminoiminomethyl)amino]phenyl]methyl]- <i>N</i> -[1-[[<i>(E)</i> -2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl]-4-[(diaminomethylene)amino]-benzamide	595
25	5-[[[1-[[<i>(E)</i> -2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl][4-[(diaminomethylene)amino]benzoyl]amino]methyl]-2-hydroxy-benzoic acid methyl ester	612
26	<i>N</i> -[1-[[<i>(E)</i> -2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl]-4-[(diaminomethylene)amino]-benzamide	448
27	4-[(aminoiminomethyl)amino]- <i>N</i> -(2-benzofuranylmethyl)- <i>N</i> -[1-[[<i>(E)</i> -2-(4-chlorophenyl)ethenyl]sulfonyl]-3-pyrrolidinyl]-benzamide	578

Example 2

4-[[[[*(E)*-2-(4-chlorophenyl)ethenyl]sulfonyl]

[1-[4-[(diaminomethylene)amino]benzoyl]-3-pyrrolidinyl]amino]methyl]-

5 benzoic acid (Compound 21)

To a solution of Compound 1C (prepared in Example 1; 0.90 g, 2.7 mMol) in CH₂Cl₂ (30 mL) was added DIPEA (0.5 mL, 2.7 mMol) followed by *p*-chlorostyrylsulfonyl chloride (0.7 g, 2.7 mMol) and stirred at rt overnight. The reaction was cooled to rt and washed sequentially with 1N KHSO₄ and brine.

10 The combined extracts were dried (Na₂SO₄), filtered and concentrated to afford Compound 2A as an orange foam: HPLC 4.31 min; 86%. Compound 2A was treated with 20 mL of TFA and stirred for 3.25 h. The solution was concentrated to afford Compound 2B as a brown oil: 2.3 g. To a solution of Compound 2B in CH₂Cl₂ (30 mL) was added DIPEA (2.0 mL, 11.5 mMol) followed by *p*-nitrobenzoyl chloride (0.4 g, 2.3 mMol) and stirred for 1h. The solution was washed sequentially with 1N HCl, H₂O, 1N NaOH, and brine, then dried (Na₂SO₄), filtered and concentrated to yield Compound 2C: 1.5 g; HPLC: 4.0 min, 75%.

20 To a suspension of Compound 2C (1.5 g, 2.5 mMol) in MeOH (30 mL) was added SnCl₂ (2.4 g, 12.7 mMol) dissolved in conc. HCl (15 mL) and the


resulting suspension heated to reflux. After 30 min at reflux, the reaction was cooled to rt, concentrated, and made basic with 1N NaOH. The mixture was extracted twice with EtOAc, and the combined extracts were dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash column chromatography (Biotage Flash 40 column, Isco UA-6 gradient pump; CH₂Cl₂:MeOH; 98:2 -> 94:6) to yield Compound **2D** as an off white solid: (0.4 g, 31%); HPLC 3.34 min, 100%; ES-MS 595 (MH⁺).

A solution of Compound **2D** (0.4 g, 0.8 mMol) and *N,N*-bis-*tert*-butoxycarbonylthiourea (*Syn. Comm.* **1993**, 23(10), 1443; 0.3 g, 1.2 mMol) in 15 mL of DMF was treated with TEA (0.4 mL, 2.6 mMol) followed by HgCl₂ (1.8 mMol, 0.5 g) and the reaction was stirred overnight. The black suspension was diluted with EtOAc (150 mL) and filtered through Celite. The filtrate was washed sequentially with H₂O and brine, filtered, dried (Na₂SO₄), and concentrated. The residue was purified by flash column chromatography (Biotage Flash 40 column, Isco UA-6 gradient pump; CH₂Cl₂:MeOH; 100% -> 96:4) to yield Compound **2E** as a white solid: (0.5 g, 73%); HPLC 4.51 min, 100%.

To a solution of Compound **2E** (0.4 g, 0.5 mMol) in 1,4-dioxane (18 mL) was added a solution of LiOH·H₂O (58 mg, 1.4 mMol) in H₂O (2 mL) and stirred for 48 hr. The solution was concentrated, acidified with 1N KHSO₄, and extracted twice with EtOAc. The combined extracts were washed with brine, dried over Na₂SO₄, filtered through Celite and concentrated. The residue was purified by flash column chromatography (Biotage Flash 40 column, Isco UA-6 gradient pump; CH₂Cl₂:MeOH; 100% -> 94:6) to yield Compound **2F** as a white powder: (0.2 g, 33%); HPLC: 4.31 min, 100%.

A solution of Compound **2F** (63 mg, 0.08 mMol) in 2 mL of 100% TFA was stirred for 45 min. The solution was concentrated to yield a clear oil which was triturated twice with Et₂O and dried under vacuum at rt to yield Compound **21** as a white solid: (43 mg, 77%); HPLC: 3.0 min 94%; ES-MS 582 (MH⁺); ¹H NMR (DMSO-d₆) δ 1.75-2.13 (m, 2H), 3.12-3.79 (overlapping m, 4H), 4.31-4.68

(overlapping m, 3H), 7.08-8.11 (overlapping m, 18H), 10.0 (broad s, 1H).



- Using the procedure of Example 2 and the appropriate reagents and starting materials known to those skilled in the art, other compounds of the present invention may be prepared including, but not limited to:

Cpd	Name	ES-MS m/z (MH ⁺)
16	2-(4-chlorophenyl)-N-[1-[4-[(diaminomethylene)amino]benzoyl]-3-pyrrolidinyl]-N-(phenylmethyl)-(E)-ethenesulfonamide	538
17	2-(4-chlorophenyl)-N-[1-[4-[(diaminomethylene)amino]benzoyl]-3-pyrrolidinyl]-N-[[4-(phenylmethoxy)phenyl]methyl]-(E)-ethenesulfonamide	645

18	2-(4-chlorophenyl)-N-[1-[4- [(diaminomethylene)amino]benzoyl]-3-pyrrolidinyl]- N-[[4-(dimethylamino)phenyl]methyl]-(E)- ethenesulfonamide	582
19	2-(4-chlorophenyl)-N-[1-[4- [(diaminomethylene)amino]benzoyl]-3-pyrrolidinyl]- N-[[4-methoxyphenyl]methyl]-(E)- ethenesulfonamide	569
20	2-(4-chlorophenyl)-N-[1-[4- [(diaminomethylene)amino]benzoyl]-3-pyrrolidinyl]- N-(3-phenylpropyl)-(E)-ethenesulfonamide	567
22	4-[[[(E)-2-(4-chlorophenyl)ethenyl]sulfonyl][1-[4- [(diaminomethylene)amino]benzoyl]-3- pyrrolidinyl]amino]methyl]-benzoic acid methyl ester	596
23	N-[1-[4-[(diaminomethylene)amino]benzoyl]-3- pyrrolidinyl]-7-methoxy-N-(phenylmethyl)-2- naphthalenesulfonamide	558

Biological Examples

The utility of the compounds of the present invention as serine protease or dual-serine protease inhibitors and, particularly, as Factor Xa or tryptase inhibitors useful as agents for the treatment of serine protease or dual-serine protease mediated disorders can be determined according to the procedures described herein.

Enzyme-Catalyzed Hydrolysis Assays

Factor Xa Inhibition

Enzyme-catalyzed hydrolysis rates were measured spectrophotometrically using commercial human Factor Xa (American Diagnostica), the chromogenic substrate (MeO-CO-D-CHG-Gly-Arg-pNa (American Diagnostica); in aqueous buffer (50 mM Trisma Base, 0.1% Tween 80, pH 8.4); and a microplate reader (Molecular Devices). Changes in absorbance at 405 nM were monitored using the software program Softmax (Molecular Devices), upon addition of enzyme, with and without inhibitor present at 37°C for 30 minutes. The IC₅₀ values were determined by fixing the enzyme and substrate concentrations (5.7 nM Factor Xa, 500 μM Factor Xa substrate) and varying the inhibitor concentration. Percent inhibition was calculated by comparing the initial reaction slopes of the

without inhibitor samples to those with inhibitor. Inhibition constants (K_i) were determined by fixing the enzyme concentrations (5.7 nM Factor Xa) and inhibitor concentrations and varying the substrate concentrations (30 – 700 μ M Factor Xa substrate. Michaelis-Menton kinetics were applied to the initial
5 reaction slopes using the program Kcat (Bio Metallics Inc.).

Table 1 summarizes assay results for Factor Xa inhibition for certain compounds of the present invention:

10

Table 1

Cpd	Factor Xa K_i (μ M)
1	1.6
2	0.3
3	0.4
4	7.7
5	0.2
6	0.6
7	0.3
8	1.1
9	0.3
10	0.2
11	0.6
12	0.4
13	0.2
14	0.3
15	0.6
17	4.4
22	7.1
23	21
27	0.1

Tryptase Inhibition

The rate of increase in absorbance at 405 nM due to hydrolysis of synthetic chromogenic peptide substrates ([S]: 500 μ M N-p-Tosyl-GLY-PRO-
15 LYS-pNA; Sigma T-6140) is measured in the presence and absence of inhibitors (I) with a microplate reader at 37°C. The enzyme reaction is started

by the addition of enzyme ([E]: 1.0 nM human Lung Tryptase; Cortex Biochem CP3033). Data is collected over a period of 30 min. and the initial rate of substrate hydrolysis (V_o (mOD/min)) is calculated. Inhibition is calculated by comparing to wells containing no inhibitor (vehicle) and IC_{50} s are determined
5 using a four parameter fit logistics model.

Table 2 summarizes assay results for tryptase inhibition for certain compounds of the present invention:

10

Table 2	
Cpd	Tryptase IC_{50} (μ M)
2	0.9
24	4.0
25	0.7
26	0.6


15

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.


CLAIMS

What is claimed is:

1. A compound selected from the group consisting of Formula (I) and
 5 Formula (II):



Formula (I)



Formula (II)

wherein

- R₁ is selected from the group consisting of hydrogen, C₁₋₈alkyl, C₃₋₇cycloalkyl, aryl, aryl(C₁₋₈)alkyl, aryl(C₂₋₈)alkenyl, heteroaryl(C₁₋₈)alkyl, heteroaryl(C₂₋₈)alkenyl and R₄C(O)CH₂-; wherein aryl and heteroaryl are
 10 optionally substituted with one to two substituents independently selected from R₄;

- R₂ is selected from the group consisting of hydrogen, hydroxy, C₁₋₈alkoxy, aryloxy and aryl(C₁₋₈)alkoxy; with the proviso that R₂ is bonded to the heterocyclyl ring by a single bond; alternatively, R₂ is oxo; with the proviso
 15 that R₂ is bonded to the heterocyclyl ring by a double bond;

R₃ is selected from the group consisting of aryl, aryl(C₁₋₈)alkyl, aryl(C₂₋₈)alkenyl, heteroaryl(C₁₋₈)alkyl, heteroaryl(C₂₋₈)alkenyl; wherein aryl and heteroaryl are optionally substituted with one to three substituents independently selected from the group consisting of halogen, C₁₋₈alkyl, C₁₋₈alkoxy, amino, (C₁₋₄alkyl)amino, di(C₁₋₄alkyl)amino, trihalo(C₁₋₈)alkyl and trihalo(C₁₋₈)alkoxy;

R₄ is selected from the group consisting of hydroxy, amino, C₁₋₈alkyl, (C₁₋₄alkyl)amino, di(C₁₋₄alkyl)amino, C₁₋₈alkoxy, carboxy, carboxy(C₁₋₈)alkyl, carboxy(C₁₋₈)alkoxy, (carboxy)amino, (carboxy(C₁₋₄)alkyl)amino, (carboxyaryl)amino, (carboxyaryl(C₁₋₄)alkyl)amino, (carboxy(C₁₋₄)alkylaryl)amino, aryloxy, aryl(C₁₋₈)alkoxy, (aryl)amino, (aryl(C₁₋₄)alkyl)amino, (C₁₋₄alkylaryl)amino, (arylcarboxy)amino, di(aryl)amino, di(aryl(C₁₋₄)alkyl)amino, C₁₋₈alkoxycarbonyl, C₁₋₈alkoxycarbonyl(C₁₋₈)alkoxy, aminocarbonyl, (C₁₋₈alkyl)aminocarbonyl, (carboxy(C₁₋₈)alkyl)aminocarbonyl, (C₁₋₈alkoxycarbonyl(C₁₋₈)alkyl)aminocarbonyl and guanidino; and,

G is selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄alkyl, C₁₋₈alkoxy, aryl, aryloxy, aryl(C₁₋₈)alkyl, aryl(C₁₋₈)alkoxy, amino, carboxy, alkylaminocarbonyl, alkylcarbonylamino, trihalo(C₁₋₈)alkyl and trihalo(C₁₋₈)alkoxy;

and pharmaceutically acceptable salts thereof.

2. The compound of claim 1 wherein R₁ is selected from the group consisting of hydrogen, aryl(C₁₋₈)alkyl and heteroaryl(C₁₋₈)alkyl, wherein the aryl and heteroaryl portion of arylalkyl and heteroarylalkyl are optionally substituted with a substituent selected from R₄.


3. The compound of claim 1 wherein R₁ is selected from the group consisting of hydrogen, benzyl, phenethyl, phenylpropyl and benzofurylmethyl, wherein phenyl, the phenyl portion of benzyl and the benzofuryl portion of benzofurylmethyl are optionally substituted with a

substituent selected from R₄.

4. The compound of claim 1 wherein R₁ is selected from the group consisting of hydrogen, benzyl, phenylpropyl and benzofurylmethyl, wherein phenyl, the phenyl portion of benzyl and the benzofuryl portion of benzofurylmethyl are optionally substituted with a substituent selected from R₄.
5. The compound of claim 1 wherein R₂ is hydrogen.
6. The compound of claim 1 wherein R₃ is aryl(C₂₋₈)alkenyl, wherein aryl is optionally substituted with one to three substituents independently selected from halogen.
7. The compound of claim 1 wherein R₃ is a substituent independently selected from the group consisting of phenethenylene and phenylpropenylene, wherein phenyl is optionally substituted with one to three substituents independently selected from the group consisting of chlorine and fluorine.
8. The compound of claim 1 wherein R₃ is phenethenylene, wherein phenyl is substituted with one to three substituents selected from chlorine.
9. The compound of claim 1 wherein R₄ is selected from the group consisting of hydroxy, di(C₁₋₄alkyl)amino, C₁₋₈alkoxy, carboxy, carboxy(C₁₋₈)alkoxy, aryl(C₁₋₈)alkoxy, C₁₋₈alkoxycarbonyl, C₁₋₈alkoxycarbonyl(C₁₋₈)alkoxy, aminocarbonyl, (C₁₋₈alkyl)aminocarbonyl, (carboxy(C₁₋₈)alkyl)aminocarbonyl and C₁₋₈alkoxycarbonyl(C₁₋₈)alkyl)aminocarbonyl.
10. The compound of claim 1 wherein R₄ is selected from the group consisting of hydroxy, carboxy, carboxy(C₁₋₈)alkoxy, C₁₋₈alkoxycarbonyl, C₁₋₈alkoxycarbonyl(C₁₋₈)alkoxy, aminocarbonyl,

(carboxy(C₁₋₈)alkyl)aminocarbonyl and
C₁₋₈alkoxycarbonyl(C₁₋₈)alkyl)aminocarbonyl.

11. The compound of claim 1 wherein R₄ is selected from the group consisting of hydroxy, carboxy, carboxymethoxy, methoxycarbonyl, aminocarbonyl, (carboxymethylene)aminocarbonyl and methoxycarbonylmethylene)aminocarbonyl.
12. The compound of claim 1 wherein G is selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄alkyl, C₁₋₈alkoxy, aryl, aryloxy, aryl(C₁₋₈)alkyl, aryl(C₁₋₈)alkoxy, amino and trihalo(C₁₋₈)alkyl.
13. The compound of claim 1 wherein G is hydrogen.
14. The compound of claim 1 of the formula:




wherein R₄ is selected from the group consisting of

R ₄
4-OH;
4-CO ₂ CH ₃ ;
3-CO ₂ CH ₃ ;
4-CO ₂ H;
3-CO ₂ H;
3-OH;
3-OCH ₂ CO ₂ CH ₃ ;
4-CONH ₂ ;
4-CONHCH ₂ CO ₂ CH ₃ ;
3-OCH ₂ CO ₂ H;

4-CONHCH₂CO₂H;
 3-CONHCH₂CO₂CH₃;
 3-CONHCH₂CO₂H;
 3-CONH₂;
 4-NHC(=NH)NH₂; and,
 3-CO₂CH₃-4-OH;

and pharmaceutically acceptable salts thereof.

15. The compound of claim 1 of the formula:




wherein R₁ is:

R₁
 PhCH₂;
 H; and,
 2-benzofuranylCH₂;










5 and pharmaceutically acceptable salts thereof.

16. The compound of claim 1 of the formula:



wherein R₁ and R₃ are dependently selected from the group consisting of

<u>R₁</u>	<u>R₃</u>
PhCH ₂	4-ClPh(CH ₂) ₂ ;
4-(PhCH ₂ O)PhCH ₂	4-ClPh(CH ₂) ₂ ;



and pharmaceutically acceptable salts thereof.

17. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.
- 5 18. A pharmaceutical composition made by mixing a compound of claim 1 and a pharmaceutically acceptable carrier.
- 10 19. A process for preparing a pharmaceutical composition comprising mixing a compound of claim 1 and a pharmaceutically acceptable carrier.
- 15 20. A method for treating a serine protease or dual-serine protease mediated disorder in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a compound of claim 1.
- 20 21. The method of claim 20 wherein the disorder is mediated by selective inhibition of a serine protease.
22. The method of claim 21 wherein the serine protease is selected from the group consisting of Factor Xa and tryptase.
- 25 23. The method of claim 20 wherein the disorder is mediated by dual inhibition of at least two serine proteases.
24. The method of claim 23 wherein the serine protease is selected from the group consisting of at least Factor Xa and tryptase.

25. The method of claim 20 wherein the serine protease or dual-serine protease mediated disorder is selected from the group consisting of thrombotic disorders, arterial thrombosis, venous thrombosis, restenosis, hypertension, heart failure, arrhythmia, myocardial infarction, acute myocardial infarction, reocclusion following thrombolytic therapy, reocclusion following angioplasty, inflammation, angina, unstable angina, stroke, atherosclerosis, ischemic conditions, neurodegenerative disorders (associated with thrombotic or ischemic conditions), asthma and inflammatory bowel syndrome.
- 5
- 10
26. The method of claim 20 wherein the therapeutically effective amount of the compound of claim 1 is from about 0.001 mg/kg/day to about 300 mg/kg/day.

INTERNATIONAL SEARCH REPORT

I Application No

PCT/US 02/06475

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 C07D207/48 C07D207/14 A61K31/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 57934 A (DU PONT MERCK PHARMA) 23 December 1998 (1998-12-23) page 1, line 1-11; claim 1	1-26
A	US 5 770 600 A (ARDECKY ROBERT JOHN ET AL) 23 June 1998 (1998-06-23) column 1, line 15-24; claim 1	1-26
A	EP 0 623 595 A (SQUIBB BRISTOL MYERS CO) 9 November 1994 (1994-11-09) page 26, line 24-54; claim 1	1-26

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the International search

19 July 2002

Date of mailing of the International search report

26/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Samsam Bakhtiary, M

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.1

Although claims 20-26 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 02/06475

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: —
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/US 02/06475

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9857934	A	23-12-1998	AU 7976998 A EP 0993448 A1 WO 9857934 A1	04-01-1999 19-04-2000 23-12-1998
US 5770600	A	23-06-1998	US 5658939 A US 5681844 A US 5776927 A AT 195947 T AU 2360995 A DE 69518648 D1 EP 0765340 A1 JP 9512022 T WO 9528420 A1 US 5646165 A	19-08-1997 28-10-1997 07-07-1998 15-09-2000 10-11-1995 05-10-2000 02-04-1997 02-12-1997 26-10-1995 08-07-1997
EP 0623595	A	09-11-1994	AU 6183494 A CA 2122397 A1 EP 0623595 A1 JP 6329627 A US 5583146 A US 5741792 A US 5741799 A	10-11-1994 04-11-1994 09-11-1994 29-11-1994 10-12-1996 21-04-1998 21-04-1998